

The Effectiveness of a Pre-Exercise Performance Drink (PRX) on Indices of Maximal Cardiorespiratory Fitness

Allyn Byars¹, Mike Greenwood², Lori Greenwood², & Warren K. Simpson³

Department of Physical Therapy, Angelo State University, San Angelo, TX¹; Exercise & Sport Nutrition Lab, Baylor University, Waco, TX², Department of Fitness and Sport Studies, Hardin-Simmons University, Abilene, TX³. Address Correspondence to Allyn.Byars@angelo.edu

Received May 5, 2006/Accepted June 6, 2006

ABSTRACT

This study examined the effectiveness of a pre-exercise drink (PRX) called EM-PACT on indices of maximal cardiorespiratory fitness. Twenty-four males (n=12) and females (n=12) ages 18-24 years (20.25 + 1.42), volunteered as subjects. Each subject performed two randomized trials of a VO_{2max} treadmill test within a week of each other. Subjects in this randomized, placebo controlled, counter balanced, crossover design, ingested either a placebo (water) or PRX 20 minutes before each exercise bout. VO_{2max} and time to exhaustion (Time) during graded exercise testing were evaluated. Using paired samples t-tests, significantly greater mean values were found in VO_{2max} and Time for the PRX trial compared to the placebo trial ($p < .05$). Results indicate that indices of cardiorespiratory fitness; specifically VO_{2max} and Time are enhanced by ingestion of PRX prior to exercise testing. The combined results of this investigation may provide meaningful practical applications for coaches and athletes alike regarding ergogenic hydration options. *Journal of the International Society of Sports Nutrition. 3(1):56-59, 2006*

Key Words: sports nutrition, ergogenic aids, VO_{2max} , aerobic performance, sport drink

INTRODUCTION

The human body's maximal ability to use or consume oxygen for aerobic metabolism during exercise, better known as VO_{2max} , is an important predictor of athletic performance in endurance activities¹. In addition, ventilatory threshold or the nonlinear increase in ventilation that coincides with the beginning of glycolysis (anaerobic threshold) for energy and the onset of blood lactate is considered by many exercise physiologists to be even a better indication of an endurance athlete's capacity for aerobic power². This is particularly true when examining the metabolic demands of middle distance runners and other similar athletes in their respective sports. Either way, the ability of an individual to reduce or tolerate more lactate production or the metabolic end product caused by the excessive metabolism of carbohydrates (CHO) is an important factor in the performance of endurance athletes as well as other sports that rely heavily upon aerobic metabolic pathways³.

Previously, research has demonstrated that CHO ingestion during aerobic exercise can improve performance during exercise sessions lasting longer than 90 minutes performed at intensities greater than 70% VO_{2max} by preventing a decline in blood glucose concentration and facilitating glucose oxidation late, whereas the timing and type of CHO ingestion following exercise influences muscle glycogen restoration^{4,5}. This information is especially important for endurance athletes since CHO type and blood glucose response is important in order to optimize CHO intake either pre or post exercise.

For example, CHO ingestion immediately prior to exercise has been reported to have a negative effect on exercise performance⁶. If an athlete consumes carbohydrate-rich foods or sport drinks within 60 minutes of the beginning of an endurance exercise performance, the glucose from the ingested food or drink enters the circulation within minutes of ingestion. The subsequent rise in blood glucose concentration causes the release of the hormone insulin, which assists in clearing glucose from the

circulation. A peak in insulin concentration in the blood occurs at the time exercise begins. Consequently glucose uptake by the muscles reaches an abnormally high rate during the exercise performance. Therefore, the consumption of simple CHO, which are digested and absorbed quickly, can be detrimental to exercise performance⁶.

This high rate of clearance glucose from the blood can potentially cause hypoglycemia which in turn can produce symptoms of acute fatigue. In summary, consuming high-glycemic CHO immediately before exercising causes blood glucose to rise rapidly (glycemic response) which may trigger excessive insulin release (insulinemic response)^{7,8}.

In contrast, consuming low glycemic carbohydrate-rich foods (starch with high amylose content or moderate glycemic CHO with high dietary fiber content) in the immediate 45-60 minute pre-exercise period allows for slower glucose absorption, reducing the potential for rebound glycemic response. Typically, the optimal forms of CHO have been combinations of glucose, fructose, sucrose, and maltodextrins with or without protein or amino acids and it has been further suggested that the glycemic index of food may be a key determining factor for when food is ingested relative to exercise participation⁹⁻¹⁴.

Currently there are many sport drinks that help the body replenish CHO levels during exercise including pre-exercise formulas whose purpose is to promote the sparing of CHO by facilitating fat substrate utilization during exercise. EM-PACT (Mannatech, Inc.) is an energy and endurance pre-exercise drink (PRX) purported to increase oxygen consumption, reduce lactate production, and improve fat utilization during aerobic activity. Although anecdotal and case study evidence exists giving merit to these claims, little or no laboratory evidence has been available to support the usage of the potentially performance enhancing product for aerobic or endurance performance.

The purpose of this study was to examine the effectiveness of a pre-exercise sport drink (PRX) on indices of maximal cardiorespiratory fitness. Specifically, VO₂max and Time to Exhaustion (Time) during graded exercise testing were evaluated.

METHODS

Subjects. In this investigation, twelve male and twelve female college students (n = 24), ages 18-24

years (20.25 ± 1.42), volunteered as subjects. Subjects signed university-approved informed consent statements in compliance with the institution's research review board on the campus in which the study was conducted.

Study Design. Subjects involved in this study were asked to submit to "two" maximal oxygen consumption tests (VO₂max) within a week of each other with at least 48 hours between trials. Subjects were required to perform each maximal effort exercise test on a motor-driven treadmill. In addition, expired lung gases were examined for the purpose of determining the amount of oxygen used during exercise for VO₂max. Expired lung gases were collected by sampling air exhaled from the mouth into a mouthpiece connected to sampling hoses and gas analyzers (CPX system, Medgraphics, Maple Grove, MN). The exercise intensity began at a low level and was advanced every three minutes by increasing the speed and incline of the treadmill belt using Bruce protocol¹⁵. During the test, heart rate and time were measured continuously while blood pressure and ratings of perceived exertion were measured toward the end of each three minute stage. VO₂max was considered to have been achieved if the subject met at least two of the following criteria: 1) an RER equal to or greater than 1.15 2) plateau of VO₂ during the last stage of exercise 3) maximal heart rate within ± 10 beats per minutes of predicted values.

Prior to test participation, subjects were asked to adhere to the following pre-test instructions: 1) wear comfortable, loose-fitting clothing 2) drink plenty of fluids over the 24-hour period preceding the test 3) avoid food, tobacco, alcohol, and caffeine for 3 hours prior to taking the test 4) avoid exercise or strenuous physical activity the day of the test and 5) get an adequate amount of sleep (6 to 8 hours) the night before the test¹⁵.

Testing Procedures. Each subject arrived thirty minutes prior to each exercise trial and was given either the recommended dosage (1 Tablespoon/18 g per 8 ounces/.24 L water) of PRX or a placebo (citrus flavored water) twenty minutes prior to test participation. Administration of PRX and placebo trials were randomized with half of the subjects ingesting the placebo during the first trial and PRX during their second trial with the order reversed for the remaining subjects. Total participation time for each test was approximately 1 hour. The PRX supplement (EM-PACT) was provided from Mannatech, Inc., Coppell, TX in sealed bottles and an


open-label format with an unbiased assistant (not involved in the testing or data analysis) mixing the PRX and providing it to the subjects. Both the PRX and water placebo were provided by an assistant blinding both subjects and investigators as to the order in which water placebo or PRX was ingested. At the end of the study investigators were provided information as to the order in which the subjects were provided either the PRX or water placebo. EM-PACT is a citrus flavored 7.2% carbohydrate-electrolyte concentration energy and endurance pre-exercise drink containing a proprietary blend of the following ingredients (Total 18 g/dose): Energy and Endurance Complex Fructose (14.58 g), citric acid (1.55 g), silicon dioxide (.54), medium chain triglycerides (.25 g), creatine monohydrate (.19 g), calcium citrate (.16 g), magnesium aspartate (.13 g), magnesium succinate (.13 g), potassium aspartate (.13 g), potassium succinate (.13 g), lemon oil powder (.09 g), choline bitartrate (.05 g), L-carnitine (.05 g), Ambrotose (Aloe vera extract: Manapol), Powder (.03) and lecithin (.01 g).

Statistical Analysis. Data were analyzed with paired sample t-tests. VO_{2max} ($ml \cdot kg^{-1} \cdot min^{-1}$) and Time (minutes) were the specific dependent variables examined in this study. In addition measures of effect size were also calculated. An alpha level of .05 was used in determining statistical significance. Statistical analyses were performed using SPSS for Windows version 12.0 statistical package (SPSS, Inc., Chicago, IL) ¹⁶. Data are presented as means \pm SE for water and PRX trials.

RESULTS

Initial results indicated significantly greater mean values in VO_{2max} ($ml \cdot kg^{-1} \cdot min^{-1}$) for the PRX trial compared to the placebo (water) trial $t(23)=5.82$, $p<.001$. It was also found that the PRX trial exhibited significantly higher mean values in time to exhaustion (minutes) when compared to the placebo trial $t(23)=4.47$, $p<.001$. Overall differences in the various parameters are depicted in FIGURE 1. Changes in mean values among the reported variables are displayed in TABLE 1.

Figure 1



DISCUSSION

The main findings of this study were that indices of cardiorespiratory fitness, specifically VO_{2max} , and time to exhaustion were significantly ($p<.05$) enhanced by ingestion of PRX prior to graded exercise testing. In particular, overall increases were observed in VO_{2max} (15.5%) and time to exhaustion (8.7%). The results of this study also support the use of the PRX as examined in this investigation in tests of aerobic power as well as support earlier reports of ingesting a PRX consisting of low glycemic sugars (5-8%) before exercise ⁹⁻¹⁴.

Improvement of time to exhaustion claims also could possibly be substantiated as the data of this investigation support a recent study in which a mixture of CHO and medium-chain triglycerides (MCTs) resulted in increased aerobic function as marked by increases in length of time trials to exhaustion ¹⁷. Marketing for the sports enthusiast hypes MCTs as fat burners, energy sources, glycogen spasers, and muscle builders. Although MCTs do not inhibit gastric emptying as does common fat, conflicting research supports the efficacy of using MCTs solely as a means of improving fat oxidation during exercise ^{18,19} and because of its minimal contribution to the formula, the individual contribution of MCTs in improving performance is highly unlikely. However, subsequent research investigating possible metabolic and ergogenic effects of combining MCTs with CHO may hold promise. It was concluded that, during bouts of exercise requiring aerobic power, the combined results of this investigation provide meaningful practical applications for coaches and athletes alike regarding possible alternative hydration options.

Future research is warranted investigating blood lactate levels, fuel substrate utilization, gender differences, fitness levels, comparisons with other products, as well as use under various environmental and competitive conditions. In addition, further research is needed to determine if other ingredients or combinations of ingredients included in the mixture of this PRX may have influenced the results. For example, creatine monohydrate is included in this mixture, however; its amount (.15 g/dose) is probably negligible in producing any ergogenic benefit due to the limited amount and timing of the ingestion period and lack of any prior loading of creatine monohydrate by the subjects. Also, L-carnitine (.04 g/dose) has been demonstrated to have an ergogenic effect (mixed reports) however; as with the creatine the amount and the timing are probably prohibitive

concerning any benefit³. Although the results of this study favor using this particular PRX, caution should be taken regarding the findings since it is difficult to provide a feasible scientific rationale why any significant findings occurred based on the content of the product.

To the author's knowledge, no previous research has shown similar significant acute results utilizing a proprietary blend of ingredients primarily designed for use as a concentrated sports drink. In effort to substantiate or refute the efficacy of this product noted in this study, additional studies are most certainly warranted.

ACKNOWLEDGEMENTS

Gratitude is expressed to the subjects that participated in this study as well as to each of the laboratory assistants who were instrumental in the collection of the data. This study was funded by a product grant from Mannatech Inc. (Coppell, TX). The researchers in this study independently collected, analyzed, and interpreted the results from this study and have no financial interests in the results of this study. Dissemination of the results in this study does not constitute endorsement by the researchers or their institutional affiliations.

Table 1. Mean changes in various parameters

Variable	Water		95% C.I.	PRX		95% C.I.	% Increase	Effect Size
	M	SD		M	SD			
VO₂max (ml·kg⁻¹·min⁻¹) (n=24)	31.95 ± 6.55	29.18 - 34.72	36.89 ± 7.16	33.86 - 39.91	15.5*	1.19		
Time (minutes) (n=24)	9.34 ± 2.24	8.40 - 10.29	10.14 ± 2.22	9.21 - 11.09	8.7*	1.29		

REFERENCES

- National Strength and Conditioning Association. *Essentials of strength and conditioning* (2nd edition) 2000; Champaign, IL: Human Kinetics.
- Nieman, D.C. *Exercise testing and prescription: a health-related approach* (5th edition). 2002; Boston, MA: McGraw-Hill.
- McArdle, W.D., Katch, F.I., and Katch, V.L. *Exercise physiology: energy, nutrition, and human performance* (5th edition). 2001; Baltimore, MD: Lippincott, William, and Wilkins.
- Sherman WM, Jacobs KA and Leenders N. Carbohydrate metabolism during endurance exercise. In: *Overtraining in Sport*, edited by R. Kreider AF, & M. O'Toole. Champaign: Human Kinetics, 1998; p. 289-293, 300-302.
- Zachwieja JJ, Costill DL and Fink WJ. Carbohydrate ingestion during exercise: effects on muscle glycogen resynthesis after exercise. *Int J Sport Nutr* 1993; 3:418-430.
- Brown, S.P., Miller, W.C., and Eason, J. *Exercise physiology: basis of human movement in health and disease*. 2006; Baltimore, MD: Lippincott, William, and Wilkins.
- Brand-Miller, J, et al. *The G.I. factor: the glycemic index solution*. 1996; Sydney, Australia: Hodder and Stoughton.
- Fontvieille, A.M., et al. *The use of low glycemic index foods improves metabolic control of diabetes patients in a 10 week study*. *Diabet Med* 1992; ; 9: 444.
- Costill DL, Sherman WM, Fink WJ, Maresh C, Witten M and Miller JM. The role of dietary carbohydrates in muscle glycogen resynthesis after strenuous running. *Am J Clin Nutr* 1981; 34: 1831-1836.
- Blom P, A. Hostmark, O.Vaage, K. Kardel, and S. Maehlum. Effects of different post-exercise sugar diets on the rate of muscle glycogen synthesis. *Med Sci Sports Exerc* 1987; 9: 491-496.
- Burke LGC, and M. Hargreaves. Muscle glycogen storage after prolonged exercise: Effect of the glycemic index of carbohydrate feedings. *J Appl Physiol* 1993; 75:1019-1023.
- Chandler RM, Byrne HK, Patterson JG and Ivy JL. Dietary supplements affect the anabolic hormones after weight-training exercise. *J Appl Physiol* 1994; 76:839-845.
- Ivy JL. Muscle glycogen synthesis before and after exercise. *Sports Med* 1991;11: 6-19.
- Ivy JL. Glycogen resynthesis after exercise: effect of carbohydrate intake. *Int J Sports Med* 1998; 19 Suppl 2: S142-145.
- American College of Sports Medicine. *Guidelines for exercise testing and prescription* (6th edition). 2000; Philadelphia, PA: Lippincott, Williams, and Wilkins.
- SPSS *Statistical package for the social sciences*. [software version 11.5], 2003; Chicago, IL: SPSS.
- Beckers, E.J., et al. *Gastric emptying of carbohydrate-medium chain triglyceride suspensions at rest*. *Int J Sports Med* 1992; 13:58.
- Jeukendrup, A.E., et al. *Fat metabolism in exercise: a review-part III: effects of nutritional interventions*. *Int J Sports Med* 1998; 19:371.
- Jeukendrup A.E., et al. *Effect of medium-chain triacylglycerol and carbohydrate ingestion during exercise on substrate utilization and subsequent cycling performance*. *Am J Clin Nutr* 1998; 67: 397.